

(Front of card)

WARNING CARD
EPZICOM® (abacavir sulfate and lamivudine) Tablets

Patients taking EPZICOM may have a serious allergic reaction (hypersensitivity reaction) that can cause death. If you get a symptom from 2 or more of the following groups while taking EPZICOM, call your doctor right away to determine if you should stop taking this medicine.

	Symptom(s)
Group 1	Fever
Group 2	Rash
Group 3	Nausea, vomiting, diarrhea, or abdominal (stomach area) pain
Group 4	Generally ill feeling, extreme tiredness, or achiness
Group 5	Shortness of breath, cough, or sore throat

Always carry this Warning Card with you to help recognize symptoms of this allergic reaction.

(Back of Card)

WARNING CARD
EPZICOM® (abacavir sulfate and lamivudine) Tablets

If you must stop treatment with EPZICOM because you have had an allergic reaction to abacavir, **NEVER** take EPZICOM or another abacavir-containing medicine (ZIAGEN® and TRIZIVIR®) again. If you take EPZICOM or another abacavir-containing medicine again after you have had an allergic reaction, **WITHIN HOURS** you may get **life-threatening symptoms** that may include **very low blood pressure** or **death**.

Please read the Medication Guide for additional information on EPZICOM.

March 2009

EPZ:2WC